

Visual Observability of the Cassiopeia A Supernova

J. A. MORGAN

*The Aerospace Corporation*¹

Abstract

It is generally believed that the explosion which gave birth to the Cassiopeia A supernova remnant resulted from core collapse of a hydrogen-deficient star. A progenitor that has lost all its hydrogen envelope and part of its helium envelope would lead to an explosion with the optical properties of a Type Ic supernova. There is evidence, if not general agreement, that Flamsteed observed the Cas A supernova as a sixth magnitude object in August, 1680. If an explosion with a typical SNIc light curve at the position and distance of Cas A attained maximum luminosity during the winter of 1679-1680, it would at that time have been poorly situated for visual observation, as its upper culmination would have taken place during daylight, while in August, between 170-200 days after peak luminosity, it would have been a sixth magnitude star.

A persistent enigma regarding the origins of the Cassiopeia A supernova remnant is the dearth of contemporary accounts of a bright new star at the time of the outburst as estimated from the remnant expansion age. Kinematical studies of the high proper-motion clouds known as fast-moving knots point to an explosion in 1672 ± 18 .² Reed et al.³ find a distance to the Cas A remnant of $d = 3.4^{+0.3}_{-0.1}$ kpc, obtained from radial velocities and proper motions of a large number of fast-moving knots. Their value agrees with that given earlier by Shklovski.⁴ At this distance, assuming normal extinction of one magnitude/kpc, a Type Ia supernova explosion should have had a peak visual magnitude of about -3, comparable to Venus. On the other hand, van den Bergh and Dodd⁵ estimate that the Cas A supernova might not have exceeded +2 magnitudes at maximum light, and, noting the frequency with which both Oriental and European astronomers failed to record nova outbursts during this period, conclude that the absence of contemporary observations is unsurprising. The only seventeenth century record of a star near the location of Cas A is a single report by Flamsteed, who observed the sixth magnitude star 3 Cassiopeiae on 1680 August 16 (Julian).⁶ This star has not been observed subsequently, and the validity of Flamsteed's observation has been contested.⁷ Setting aside the question of whether or not Flamsteed did observe a star near the present-day location of the Cas A remnant, it seems worth trying to construct a plausible sequence of events that explains both the disputed observation of 3 Cassiopeiae by Flamsteed and the lack of other contemporary reports of a new star. This note examines the possibility that Cas A was a Type Ic supernova (at least, as to light curve) that attained peak luminosity in the winter of 1679-1680, most probably in February or March, 1680. Flamsteed's observation of 3 Cassiopeiae, therefore, would have been made some six months after maximum light of the Cas A supernova. As will become apparent, the account to be presented does not tell a complete tale, and is at points necessarily conjectural.

It has long been thought that the Cas A supernova resulted from core collapse of a hydrogen-deficient star. The discovery⁸ of a compact X-ray source, interpreted as a neutron star,⁹ near the inferred expansion center of the fast-moving knots and the observation of significant ^{44}Ti γ -emission¹⁰ provide direct evidence for a core-collapse origin. Very little hydrogen emission has been observed in the fast-moving knots. If we posit a progenitor that has been stripped of its hydrogen envelope altogether, the resulting explosion would be an SNIb or SNIc supernova.¹¹

There are difficulties with assuming that Cas A supernova was of Type Ic. To produce an SNIc light curve requires a progenitor that is (1) compact and (2) of low mass. No direct evidence bearing on the radius of the Cas A progenitor at the time of the explosion is available to us, but from the remnant one might hope to learn somewhat concerning its mass. Estimates of the Cas A progenitor mass at the time of the explosion vary widely. One-dimensional hydrodynamic calculations of the explosion of helium stars that have undergone significant mass loss reproduce the light curve of an SNIc outburst with a progenitor mass between $2.3M_{\odot} - 3.6M_{\odot}$ ^{12,13} This mass range comports awkwardly with estimates of the supernova ejecta mass of order $4M_{\odot}$.^{14,15} Measured abundance ratios of nucleosynthetic products appear consistent with a mass of ca. $12M_{\odot}$,¹⁷ while a recent three-dimensional numerical study of the evolution and explosion of Cas A finds that a WN Wolf-Rayet star of mass between $4M_{\odot} - 6M_{\odot}$ best fits combined constraints posed by nucleosynthesis, ejecta mass, and compact remnant.¹⁸ Explosion of a Wolf-Rayet star of appreciable mass would presumably result in an SNIb outburst. There is also evidence that the Cas A progenitor may have had a very thin layer of hydrogen¹⁷. However, the light curve of the peculiar SNII SN1993J also fits the requirements of the scenario presented here.

Cas A lies near the Galactic plane at the low latitude of -2.1° , and extinction appears to be nonuniform across the remnant. Hurford and Fesen¹⁹ have determined from $[SII]$ line ratios that the extinction to five fast-moving knots lies in the range $A_v = 4.6 - 5.4$ within errors estimated as $\pm(0.25 - 0.45)$. Using (less certain) Balmer $H\alpha/H\beta$ ratios, they also find $A_v \leq 5.3^{+0.9}_{-0.9}$ and $\leq 6.2^{+0.9}_{-0.9}$ to two of the low proper-motion quasistationary flocculi. These values are about one magnitude greater than the value $A_v = 4.3$ previously found by Searle,²⁰ and suggest that the extinction varies across the remnant by as much as a magnitude. The distance modulus for Cas A is $12.7^{+0.2}_{-0.1}$. If we provisionally accept Flamsteed's observation of a sixth magnitude star near the location of the present-day Cas A supernova remnant, and estimate the visual extinction to that remnant to be $A_v = 5^{+0.45}_{-0.45}$, then the absolute magnitude of the supernova was approximately $M_V = -11.7^{+0.5}_{-0.4}$ in August, 1680. This value is characteristic of late > 200 day behavior of a supernova light curve, rather than an early peak value of approximately $M_V = -16$ to -18 , suggesting that the initial outburst could have taken place as much as 200 days earlier than 1680 August 16.

At any time between, roughly, New Year's 1680 and early June of that year, the (circumpolar) Cas A supernova would have had its upper culmination in the daytime sky in

the Northern hemisphere. First observation of the supernova would almost certainly have been with naked eye, despite the general use of the telescope²¹ by late seventeenth-century astronomers. A Type Ic supernova at the distance of Cas A would have been unobservable by naked eye in daytime even at peak brightness. The threshold visual magnitude for observation of a star by a human observer in the presence of Rayleigh-scattered sunlight may be estimated as described by Hughes²². The minimum detectable brightness contrast in *lux* detectable by human observers^{23,24} is converted to stellar magnitude with Russell's^{25,26} value for the stellar equivalent $1 \text{ lux} = -14.18$ visual magnitudes. The atmospheric radiative transfer code MODTRAN4^{27,28} was used to calculate the brightness of the sunlit sky at the February 14, 1680 position of the expansion center of Cas A as viewed from Greenwich for Julian dates 1680 February 2, February 12, February 22/23, and March 3, using Solar positions from Gingerich and Welther²⁸. Threshold m_V magnitudes corresponding to a just-detectable star at these dates for Cas A altitudes near upper culmination appear in Table I. The m_V values in the table are probably in error by no more than ± 0.1 magnitudes. Threshold m_V values are higher for very low Solar altitudes, samples of which also appear in the table, but the plane-parallel MODTRAN model is probably not to be relied upon for solar altitudes as low as 5° .

Table 1. Threshold V Magnitudes at location of Cas A

date	Cas A altitude	solar altitude	threshold m_V
1680 Feb 2	83.66°	22.62°	-2.4
"	65.22°	5.75°	-1.0
1680 Feb 12	83.40°	28.64°	-3.1
"	56.65°	5.04°	-0.8
1680 Feb 22	51.78°	8.07°	-1.3
1680 Feb 23	83.76°	33.14°	-2.8
1680 Mar 3	82.57°	36.66°	-3.0
"	42.71°	5.51°	-1.2

For purposes of discussion, consider SN1994I as a template for the Cas A outburst.

Table 2. Apparent V Magnitudes of SN1994I and SN1993J at distance of Cas A

template	A_v	peak m_V	$m_V(170d)$	$m_V(185d)$	$m_V(200d)$	$m_V(220d)$	$m_V(235d)$	$m_V(250d)$
SN1994I	5.0	-0.43	5.0	5.3	6.0	—	—	—
”	5.3	-0.13	5.3	5.6	6.3	—	—	—
”	6.0	0.57	5.5	5.8	6.5	—	—	—
SN1993J	5.0	0.04	—	—	4.6	5.0	5.2	5.5
”	5.3	0.34	—	—	4.9	5.3	5.5	5.8
”	6.0	1.04	—	—	5.5	6.0	6.2	6.5

Estimated apparent V magnitudes for SN1994I at the distance of Cas A for maximum light and various times thereafter appear in Table II.³⁰ If the values shown for A_v are assumed uncertain by ± 0.45 magnitudes, the apparent V magnitudes of Cas A in the table should have a net uncertainty $(-0.4, +0.5)$ from combined errors in extinction and distance modulus. The uncertainty arising solely from error in the distance modulus is $(-0.1, +0.2)$ magnitudes. Table II also contains apparent V magnitudes for SN1993J at the distance of Cas A.³¹ The SN1993J V light curve is quite similar to SN1994I. While the peak M_V of SN1993J is dimmer by about half a magnitude, its exponential tail is somewhat brighter at the same time after maximum light, and at the distance of Cas A, it attains the same visual magnitude as SN1994I as much as 60 days later. Without further adjustment of the overall brightness of the light curve, it is just possible to fit it into the scenario for Cas A.

Richmond et al. find a peak visual magnitude $M_v = -18.09^{+0.58}_{-0.58}$ for SN1994I.³⁰ Taking $M_v = -18.09$ for the peak magnitude and $A_v = 5$ the peak apparent visual magnitude of Cas A would have been $m_v(peak) = -0.43^{+0.2}_{-0.1}$. In early February 1680, the threshold visual magnitude which would have been observable by naked eye against the background of Rayleigh-scattered sunlight would have been brighter than $m_v(threshold) = -1$ for Solar altitudes above 5° .³² If the Cas A explosion occurred between the new year and June, Cassiopeia would have been unobservable by naked eye anytime near upper culmination.

The foregoing estimates show that, on this account, the position of the Cas A supernova in winter 1680 would have been such that sunlight would have precluded naked-eye

discovery anywhere near the meridian. This may go some ways toward accounting for the failure of contemporary astronomers to note any such phenomenon, but leaves unexplained why the circumpolar supernova was not observed at night. Cas A would have been visible at peak brightness (whenever that occurred) during some portion of nighttime at an appreciable altitude above the horizon. On the second of February, it should have exceeded the visible threshold shortly after sunset at an altitude of approximately 57° . At this altitude, atmospheric extinction would have been about 0.1 V magnitudes, so that Cas A could have appeared as bright as $m_V = -0.3^{+0.2}_{-0.1}$ ($A_v = 5$).

It is necessary, then, to stipulate some reason for the absence of reported observations of the supernova during the winter months apart from its unobservability during daylight. Here the story, unavoidably, becomes speculative. Apart from known factors such as the small number of astronomers active in the seventeenth century and the lack of any network for collection, dissemination, or archival of reports by non-astronomers, one might adduce the discouraging effect on observation of winter conditions in the Northern hemisphere, including the possibility of extended periods of cloudy weather. The supernova could have been dimmer at its peak, or its extinction greater, and it could have decayed somewhat more slowly, than assumed to this point: On February second, an outburst with peak $M_V \geq -17$ or extinction $A_v \geq 7$ would have appeared to an observer with $m_V(\text{peak}) \geq 1.7^{+0.2}_{-0.1}$, in the range which van den Bergh and Dodd suggest might have escaped notice by contemporary observers.⁵

Return to the observation of 3 Cassiopeiae on the night of 1680 August 16⁵. Flamsteed reported stellar magnitudes as integer rank values, so that one may suppose that 3 Cassiopeiae could have been as bright as $m_V = 5.5$. The SN1994I light curve could have faded to $m_v = 5.5^{+0.2}_{-0.1}$ by day 170 after maximum ($A_v = 5$), while SN1993J could have done so by day 200 ($A_v = 6$). Note that the rapid decay of SNeI is necessary in order to reach sixth magnitude within the requisite period of 170-200 days after peak brightness; no normal SNeII light curve is known to decay that rapidly.

Although the particular scenario presented in this note requires the Cas A supernova to have been observable at the time and location of the disputed report by Flamsteed, the explosion could just as well have occurred mid to late winter of some other year in the late seventeenth century, resulting an unprepossessing object of sixth magnitude at the upper

culmination of Cassiopeia in August of that year.

-
- ¹ Institution shown for purposes of affiliation only. Electronic address: john.a.morgan@aero.org
- ² Fesen, R. A., M. C. Hammel, J. Morse, R. A. Chevalier, K. J. Borkowski, M. A. Dopita, C. L. Gerardy, S. S. Lawrence, J. C. Raymond, and S. van den Bergh, *Astrophys. J.* **645**, p. 283 (2006)
- ³ Reed, J. E., J. J. Hester, A. C. Fabian, and P. F. Winkler, *Astrophys. J.* **440**, p. 706 (1995)
- ⁴ Shklovski, I. S., *Supernovae*, John Wiley and Sons, New York, p. 86 (1968)
- ⁵ van den Bergh, S., and W. W. Dodd, *Astrophys. J.* **162**, p. 485 (1970)
- ⁶ Ashworth, W. B., *J. Hist. Astron.* **xi**, p. 1 (1980)
- ⁷ Stephenson, F. R., and D. A. Green, *Historical Supernovae and their Remnants*, Clarendon Press, Oxon., Ch. 4 (2002)
- ⁸ Tannanbaum, H., IAU Circular 7246 (1999)
- ⁹ Chakrabarty, D., M. J. Pivovarov, L. E. Hernquist, J. S. Heyl, and R. Narayan, *Astrophys. J.* **548**, p. 800 (2001)
- ¹⁰ Iyudin, A. F., R. Diehl, H. Boemen, W. Hermsen, G. G. Lichti, D. Morris, J. Ryan, V. Schönfelder, H. Steinle, M. Varendorff, C. de Vries, and C. Winkler, *Astron. Astrophys* **440**, p. L1 (1994)
- ¹¹ The possibility that the Cas A explosion might be of Type Ic was mentioned in Fesen, R. A., M. C. Hammel, J. Morse, R. A. Chevalier, K. J. Borkowski, M. A. Dopita, C. L. Gerardy, S. S. Lawrence, J. C. Raymon, and S. van den Bergh, *Astrophys. J.* **636**, p. 859 (2006)
- ¹² Woosley, S. E., N. Langer, and T. A. Weaver, *Astrophys. J.* **448**, p. 315 (1995)
- ¹³ Ensman, L. M., and S. E. Woosley, *Astrophys. J.* **333**, p. 754 (1988)
- ¹⁴ McKee, C. F., and J. K. Truelove, *Physics Reports* **256**, p. 157 (1995)
- ¹⁵ Vink, J., J. S. Kaastra, and J. A. M. Bleeker, *Astron. Astrophys.* **307**, p. L41 (1996)
- ¹⁶ Fesen, R. A., *Astrophys. J. Supp.* **133**, p. 161 (2001)
- ¹⁷ Willingale, R., J. A. M. Bleeker, K. J. van der Heyden, J. S. Kaastra, and J. Vink, *Ast. Astrophys.* **381**, p. 1039 (2002)
- ¹⁸ Young, P. A., C. L. Gryer, A. Hungerford, D. Arnett, G. Rockefeller, F. X. Timmes, B. Voit, C. Meakin, and K. A. Eriksen, *Astrophys. J.* **640**, p. 891 (2006)

- ¹⁹ Hurford, A. P., R. A. Fesen, *Astrophys. J.* **469**, pp. 246 (1996)
- ²⁰ Searle, L, *Astrophys. J.* **168**, p. 41 (1971)
- ²¹ The seven-foot sextant used by Flamsteed to measure the position of 3 Cassiopeiae had telescopic sights, *vide.* Chapman, A., *England's Leonardo: Robert Hooke and the Seventeenth-Century Scientific Revolution*, CRC Press, London, p. 90 (2005)
- ²² Hughes, D. W., *Q. J. R. astr. Soc.* **24**, p. 246 (1983)
- ²³ Knoll, H. A., R. Tousey, and E. O. Hulbert, *J. Opt. Soc. Am.* **36**, p. 480 (1946)
- ²⁴ Hecht, S., *J. Opt. Soc. Am.* **37**, p. 59 (1946)
- ²⁵ Russell, H. N., *Astrophys. J.* **43**, p. 129 (1916)
- ²⁶ Russell, H. N., *Astrophys. J.* **45**, p. 60 (1917)
- ²⁷ Berk, A., G. P. Andersen, P. K. Acharya, J. H. Chetwynd, L. S. Bernstein, E. P. Shettle, M. W. Matthew, and S. M. Adler-Golden, *MODTRAN4 User's Manual*, Air Force Research Laboratory, Hanscom AFB, MA (1999)
- ²⁸ Kniezys F., D. C. Robertson, L W Abreu, P. Acharya, G. P. Anderson, L. S. Rothman, J. H. Chetwynd, J. E. A. Selby, E. Pl Shettle, W. O. Ballery, A. Berk, S. A. Clough, and L. S. Bernstein, *The MODTRAN 2/3 Report and LOWTRAN 7 Model*, Phillips Laboratory, Hanscom AFB, MA (1996)
- ²⁹ Gingerich, O., and B. L. Welther, *Planetary, Lunar, and Solar Postions New and Full Moons, A. D. 1650-1805*, Am. Phil. Soc., Philadelphia (1983)
- ³⁰ Richmond, M. W., S. D. Ban Dyk, W. Ho, C. Peng, Y. Paik, R. R. Treffers, A. V. Fillipenko, J. Bustamatne-Donas, M. Moeller, C. Pawellk, H. Tartara, and M. Spence, *Astronomical J.* **111**, p. 327 (1996)
- ³¹ Richmond, M. W, R. R Treffers, A. V. Fillipenko, and Y. Paik, *Astronomical J.* **112**, p. 732, (1996)
- ³² Perhaps surprisingly, scintillaton of the unresolved supernova would not have greatly altered the threshold contrast for visual detection; *vide.* Robson, J. G., *J. Opt. Soc. Am.* **56**, p. 1141 (1966)